

BIBLIOGRAFIA

- (1) Roberts LJ, Beardsworth SA, Trew G. Labor following caesarean section: current practice in the United Kingdom. *Br J Obstet Gynaecol.* 1994; 101: 153-5.
- (2) Norman P, Kostovcik S, Lanning A. Elective repeat caesarean sections: how many could be vaginal births? *Can Med Assoc J* 1993; 149:431 – 5.
- (3) Kline J, Arias F. Analysis of factors determining the selection of repeated cesarean section or trial of labor in patients with history of prior cesarean delivery. *J Reprod Med* 1993; 38:289 – 92.
- (4) Society of Obstetricians and Gynaecologists of Canada. Vaginal birth after previous cesarean birth. Clinical Practice guideline N° 68. Ottawa (ON): SOGC; 1997 (December)
- (5) National Institutes of Health. Cesarean childbirth. Bethesda (MD): NIH; 1981. Publication N° 82 – 2067.
- (6) Flamm BL, Lim OW, Jones C, Fallon D, Newman LA, Mantis JK. Vaginal birth after cesarean section: results of a multicenter study. *Am J Obstet Gynecol* 1988; 158:1079 – 84.
- (7) Socol ML, Peaceman AM. Vaginal birth after cesarean. An appraisal of fetal risk. *Obstet Gynecol* 1999; 93: 674 – 9.
- (8) McMahon MJ. Vaginal birth after cesarean. *Clin Obstet Gynecol* 1998; 41: 369 – 81.
- (9) Woolf SH, Battista RN, Angerson GM, Logan AG, Eel W. Canadian Task Force on the Periodic Health Exam. Ottawa (ON): Canada Communication Group; 1994. p. XXXVII.
- (10) Biswass A. Management of previous cesarean section. *Curr Opin Obstet Gynecol* 2003; 15: 123 – 9.
- (11) Quilligan EJ. Vaginal birth after cesarean section: 270 degrees. *J Obstet Gynecol Res* 2001; 27: 169 – 73.
- (12) Scott JR. Avoiding labor problems during vaginal birth after cesarean delivery. *Clin Obstet Gynecol* 1997; 40: 553 – 41.
- (13) Rageth JC, Juzi C, Grossenbacher H. Delivery after previous cesarean: a risk evaluation. *Obstet Gynecol* 1999; 93: 332 – 7.
- (14) Lovell R. Vaginal delivery after caesarean section: factors influencing success rate. *Aust N Z J Obstet Gynecol* 1996; 36: 4 – 8.
- (15) Weinstein D, Benshushan A, Tanos V, Zilberstein R, Rojansky N. predictive score for vaginal birth after cesarean section. *Am J Obstet Gynecol* 1996; 174: 192 – 8.
- (16) Bujold E, Gauthier R. Should we allow a trial of labor after a previous cesarean for dystocia in the second stage of labour? *Obstet Gynecol* 2001; 98: 652 – 5.
- (17) Hoskins IA, Gómez JL. Correlation between maximum cervical dilatation at cesarean delivery and subsequent vaginal birth after cesarean delivery. *Obstet Gynecol* 1997; 89: 591 – 3.
- (18) Shipp TD, Zelop CM, Repke JT, Cohen A, Caughey AB, Lieberman E. Trabajo de parto despues de cesarean previa: influencia de la indicación primaria y de la paridad. *Obstet Gynecol* 2000; 95: 913 – 6.
- (19) Khan KS, Risvi A. The partograph in the management of labor following c – section. *Int J Gynecol Obstet* 1995;50: 151 – 7.
- (20) Guleria K, Dhall GI, Dhall K. Pattern of cervical dilatation in pevious lower segment c section patients. *J Indian Med Assoc* 1997;95: 131 – 4.
- (21) Weinstein D, Benshushan A, Ezra Y. Vaginal birth after cesarean section: current opinion. *Int J Gynecol Obstet* 1996;53: 1 – 10.
- (22) ACOG. Vaginal birth after previous cesarean delivery. *ACOG Practice Bulletin* n° 5. Washington DC 1999.
- (23) Phelan JP, Korst LM, Settles DK. Uterine activity patterns in uterine rupture: a case – control study. *Obstet Gynecol* 1998;92: 394 – 7.
- (24) Kaplan B, Royburt M, Peled Y. Routine revision of uterine scar after prior cesarean section. *Acta Obstet Gynecol Scand* 1994;73: 473 – 5.
- (25) Bucklin BA. Vaginal birth after cesarean delivery. *Anesthesiology* 2003;99: 1444 – 8.
- (26) McMahon MJ, Luthier ER, Bowes WA. Comparison of a trial of labor with an elective second cesarean section. *N Engl J Med* 1996;335: 689 – 95.
- (27) Quilligan EJ. Vaginal birth after cesarean section. 270 degrees. *J Obstet Gynaecol Res* 2001;27: 169 – 73.
- (28) Rageth JC, Juzi C, Grossenbacher H. Delivery after previous cesarean: a risk evaluation. *Obstet Gynecol* 1999;93: 332 – 7.
- (29) Hill DA, Chez RA, Quinlan J. Uterinr rupture and dehiscence associated with intravaginal misoprostol cervica ripening. *J Reprod Med* 2000;45: 823 – 6.
- (30) McMahon MJ. Vaginal birth after cesarean. *Clin Obstet Gynecol* 1998;41: 369 – 81.

- (31) Yap OWS, Kim ES, Laros RK. Maternal and neonatal outcomes after uterine rupture in labor. *Am J Obstet Gynecol* 2001;184: 1576 – 81.
- (32) Flamm BL, Lim OW, Jones C. Vaginal birth after cesarean section: results of a multicenter study. *Am J Obstet Gynecol* 1988;158: 1079 – 84.
- (33) Hibbard JU, Ismail MA, Wang Y, Te C. Failed vaginal birth after cesarean section. How risky is it? *Am J Obstet Gynecol* 2001;184: 1365 – 73.
- (34) Appleton B, Targett C, Rasmussen M, Readman E, Sale F. Vaginal birth after cesarean section: an Australian multicenter study. *Aust N Z J Obstet Gynaecol* 2000;40: 87 – 91.
- (35) Shipp TA, Zelop CM, Repke JT, Cohen A. Intrapartum uterine rupture and dehiscence in patients with prior lower uterine segment vertical and transverse incisions. *Obstet Gynecol* 1993;94: 735 – 40.
- (36) Martin JN, Perry KG, Roberts WE. The case for trial of labor in the patient with prior low segment vertical cesarean incision. *Am J Obstet Gynecol* 1997; 177: 144 – 8.
- (37) Adair CD, Sanchez-Ramos L, Whitaker D, McDyer DC. Trial of labor in patients with a previous lower uterine vertical cesarean section. *Am J Obstet Gynecol* 1996; 174: 966 – 70.
- (38) Stovall TG, Shaver DC, Solomon SK. Trial of labor in previous cesarean section patients excluding classical cesarean sections. *Obstet Gynecol* 1987; 70: 713 – 7.
- (39) Shimonovitz S, Botosneano A, Hochner-Celnikier D. Successful first vaginal birth after cesarean section: a predictor of reduced risk for uterine rupture in subsequent deliveries. *J Indian Med Assoc* 2000, 2: 526 – 8.
- (40) Rosen MG, Dickinson JC, Westhoff CL. Vaginal birth after cesarean section: a meta – analysis of morbidity and mortality. *Obstet Gynecol* 1991; 77: 465 – 70.
- (41) Mozurkewich EL, Hutton EK. Elective repeat cesarean delivery versus trial of labor: a meta – analysis of the literature from 1989 to 1999. *Am J Obstet Gynecol* 2000; 183: 1187 – 97.
- (42) Kieser KE, Baskett TF. A 10 – year population – based study of uterine rupture. *Am J Obstet Gynecol* 2002; 100: 749 – 53.
- (43) Bujold E, Gauthier RJ. Neonatal morbidity associated with uterine rupture: what are the risk factors? *Am J Obstet Gynecol* 2002; 186: 311 – 4.
- (44) Chauhan SP, Martin JN, Henrichs CE, Morrison JC, Magann EF. Maternal and perinatal complications with uterine rupture in 142,075 patients who attempted vaginal birth after cesarean delivery: a review of the literature. *Am J Obstet Gynecol* 2003; 189: 408 – 17.
- (45) Hamilton EF, Bujold E, McNamara H, Gauthier R, Platt RW. Dystocia among women with symptomatic uterine rupture. *Am J Obstet Gynecol* 2001; 184: 620 – 4.
- (46) Flamm BL, Goings JR, Fuelbirth N, Fischmann E, Jones C, Hewson SA. Oxytocin during labor after previous cesarean section: results of a multicenter study. *Obstet Gynecol* 1987;70: 709 – 12)
- (47) Zelop CM, Shipp TA, Repke JT, Cohen A, Lieberman E. Uterine rupture during unduced or augmented labor in gravid women with one prior cesarean delivery. *Am J Obstet Gynecol* 1999; 181: 882 – 6.
- (48) Goetzl L, Shipp TA, Cohen A, Zelop ZM, Repke JT, Lieberman E. Oxytocin dose and the risk of uterine rupture in trial of labor after cesarean. *Obstet Gynecol* 2001; 97: 381 – 4.
- (49) Ravasia DJ, Wood SL, Pollard JK. Uterine rupture during induced trial of labor among women with previous cesarean delivery. *Am J Obstet Gynecol* 2000; 183: 1176 – 9.
- (50) Sanchez - Ramos L, Gaudier FL, Kaunitz AM. Cervical ripening and labor induction after previous cesarean delivery. *Clin Obstet Gynecol* 2000; 43: 513 – 23.
- (51) Delaney T, Young DC. Spontaneous versus induced labor after a previous cesarean delivery. *Obstet Gynecol* 2003; 102: 39 – 44.
- (52) Lyndon – Rochelle M, Holt VL, Easterling TR, Martin DP. Risk of uterine rupture during labor among women with a prior cesarean delivery. *N Engl J Med* 2001; 345: 3 – 8.
- (53) Katz VL, Farmer RM, Dean CA, Carpenter ME. Use of misoprostol for cervical ripening. *South Med J* 2000; 93: 881 – 4.
- (54) Choy – Hee L, Raynor BD. Misoprostol induction of labor among women with a history of cesarean delivery. *Am j Obstet Gynecol* 2001; 184: 1115 – 7.
- (55) Cunha M, Bugalho A, Bique C, Berstrom S. Induction of labor by vaginal misoprostol in patients with previous cesarean delivery. *Acta Obstet Gynecol Scand* 1999; 78: 653 – 4.
- (56) Plaut MM, Schwartz ML, Lubarsky SL. Uterine rupture associated with the use of misoprostol in the gravid patient with a previous cesarean section. *Am J Obstet Gynecol* 1999; 180: 1535 – 42.
- (57) Wing DA, Lovett K, Paul RH. Disruption of prior uterine incision following misoprostol for labor induction in women with previous cesarean delivery. *Obstet Gynecl* 1998; 91: 828 – 30.

- (58) Miller DA, Diaz FG, Paul RH. Vaginal birth after cesarean: a 10 – year experience. *Obstet Gynecol* 1994; 84: 255 – 8.
- (59) Asakura H Myers SA. More than one previous cesarean delivery: a 5 – year experience with 435 patients. *Obstet Gynecol* 1995; 85: 924 – 9.
- (60) Chattopadhyay SK, Sherbeeni MM, Anokute CC. Planned vaginal delivery after two previous caesarean sections. *Br J Obstet Gynaecol* 1994; 101: 498 – 500.
- (61) Bretelle f, Cravello L, Shojai R, Roger V, D'ercole C, Blanc B. Vaginal birth following two previous cesarean sections. *Eur J Obstet Gynecol Reprod Biol* 2001; 94: 23 – 6.
- (62) Phelan JP, Ahn MO, Diaz F, Brar HS, Rodriguez MH. Twice a cesarean, always a cesarean? *Obstet Gynecol* 1989; 73: 161 – 5.
- (63) Porreco RP, Meier PR. Trial of labor in patients with multiple previous cesarean sections. *J Reprod Med* 1983; 28: 770 – 2.
- (64) Farmakides G, Duvivier R, Schulman H, Schneider E, Biordi J. Vaginal birth after two or more previous cesarean sections. *Am J Obstet Gynecol* 1987; 156: 565 – 6.
- (65) Miller DA, Mullin P, Hou D, Paul RH. Vaginal birth after cesarean section in twin gestation. *Am J Obstet Gynecol* 1996;175: 194 – 8.
- (66) Strong TH, Phelan JP, Ahn MO, Sarno AP. Vaginal birth after cesarean section in the twin gestation. *Am J Obstet Gynecol* 1989; 161: 29 – 32.
- (67) Odeh M, Tarazova L, Wolfson M, Oettinger M. Evidence That women with a history of cesarean section can deliver twins safely. *Acta Obstet Gynecol Scand* 1997; 76: 663 – 6.
- (68) Myles T. Vaginal birth of twins after previous cesarean section. *J Matern Fetal Med* 2001; 10: 171 – 4.
- (69) Wax JR, Philput C, Mather J, Steinfeld JD, Ingardia CJ. Twin vaginal birth after cesarean. *Conn Med* 2000; 64: 205 – 8.
- (70) Sansregret A, Bujold E, Gauthier RJ. Twin delivery after a previous cesarean: a twelve – year experience. *J Obstet Gynaecol Can* 2003; 25: 294 – 8.
- (71) Delaney T, Young DC. Trial of labor compared to elective cesarean in twin gestations with a previous cesarean delivery. *J Obstet Gynaecol Can* 2003; 25: 289 – 92.
- (72) Hannah ME, Hannah WJ, Hewson SA. Planned cesarean section versus planned vaginal birth for breech presentation at term: a randomised multicenter trial. *Lancet* 2000; 356: 1375 – 83.
- (73) Zelop CM, Shipp TA, Repke JT, Cohen A, Lieberman E. Outcomes of trial of labor following previous cesarean delivery among women with fetuses weighing > 4000g. *Am J Obstet Gynecol* 2001; 185: 903 – 5.
- (74) Flamm BL, Goings JR. Vaginal birth after cesarean section: is suspected fetal macrosomia a contraindication? *Obstet Gynecol* 1989; 74: 694 – 7.
- (75) Phelan JP, Eglinton GS, Horenstein JM. Previous cesarean birth: trial of labor in women with macrosomic infants. *J Reprod Med* 1984; 29: 36 – 40.
- (76) Esposito MA, Menihan CA, Malee MP. Association of interpregnancy interval with uterine scar failure in labor: a case – control study. *Am J Obstet Gynecol* 2000; 183: 1180 – 3.
- (77) Shipp TA, Zelop CM, Repke JT, Cohen A, Lieberman E. Interdelivery interval and risk of symptomatic uterine rupture. *Obstet Gynecol* 2001; 97: 175 – 7.
- (78) Huang WH, Nakashima DK, Rumney PJ, Keegan KA, Chan K. Interdelivery interval and the success of vaginal birth after cesarean delivery. *Obstet Gynecol* 2002; 99: 41 – 4.
- (79) Bujold E, Mehta SH, Bujold C, Gauthier R. Interdelivery interval and uterine rupture. *Am J Obstet Gynecol* 2002; 187: 1199 – 202.
- (80) Yeh S, Huang X, Phelan JP. Post- term pregnancy after previous cesarean section. *J Reprod Med* 1984; 29: 41 – 4.
- (81) Callahan C, Chescheir N, Steiner BD. Safety and efficacy of attempted vaginal birth after cesarean beyond the estimated date of delivery. *J Reprod Med* 1999; 44: 606 – 10.
- (82) Zelop CM, Shipp TA, Repke JT, Cohen A, Lieberman E. Trial of labor after 40 weeks´ gestation in women with prior cesarean. *Obstet Gynecol* 2001; 97: 391 – 3.